

Paper II-15

IRRIGATION WITH MAGNETIZED WATER, A NOVEL TOOL FOR IMPROVING CROP PRODUCTION IN EGYPT

M. HOZAYN¹, A. AMANY. ABD EL-MONEM² AND AMIRA M.S. ABDUL QADOS³

¹*Agronomy Department, Agricultural and Biology Division, National Research Centre, Dokki, Cairo, Egypt*

²*Botany Department, Agricultural and Biology Division, National Research Centre, Dokki, Cairo, Egypt*

³*Botany Department, Princess Nora Bint Abdul Rahman University. Granada eq., Riyadh, Saudi Arabia*

ABSTRACT

Agricultural sciences take an interest not only in the common and valued crop-forming factors, but also in those less expensive, safe environmentally and generally underestimated. The technology of magnetic water has been developed and subsequently used widely in the field of agriculture in many countries such as Australia, USA, China and Japan. Desperate its importance, it is not yet explored in Egypt. Therefore, the present work was carried out to study the response of some food crops using magnetized water for irrigation purpose under green house condition. Monocotyledonous such as wheat and flax and dicotyledonous such as chick-pea and lentil plants were selected for the present study. Based on the results of our experiments revealed all the plants which were irrigated with magnetic water exhibited a remarkable increase in vegetative growth and biochemical constitutions. Further the results indicated that the number of protein bands got increased in plants treated with magnetized water when compared to untreated control plants. Moreover, the magnetized water treatment increased yield and yield component traits of all crops. Over two seasons, the increases in seeds yield/plant in monocotyledonous crops reached to 9.10% and 31.3% for flax and wheat, respectively and in dicotyledonous crops reached to 24.9% and 38.5%, for lentil and check pea, respectively compared with crops irrigated with tap water. It appears that the preliminary study on utilization of magnetized water can led to improve quantity and quality of crop production under Egyptian condition.

INTRODUCTION

Till 1980 a little were known about how the magnetic field (MF) can stimulate plant growth or even prevent it. Wojcik (1995) reported that in the beginning of 1980s Japanese called Fujio Shimazaki working in Shimazaki Seed Company was the first who reported that stationary magnetic fields can improve the germination of seeds and speed up the growth of plants. It has been reported that the magnetic field influenced the maximal seed germination and yield potential (Aladjadjiyan, 2002). The reason of this effect can be searched in the presence of paramagnetic properties in chloroplast which can cause an acceleration of seeds metabolism by magnetic treatment (Aladjadjiyan and Ylieve, 2003). MF increased yield parameters of crops like cereal, sunflower and soybean (Özalpan *et al.*, 1999; Yurttas *et al.*, 1999 and Oldacay, 2002). On the other hand, it has been recorded that MF affected various characteristics of plants like germination of seeds, seedlings growth and reproduction including the growth of meristematic cells and chlorophyll development (Namba *et al.*, 1995; Atak *et al.*, 1997 and Reina *et al.*, 2001).

The effect of magnetic field on the productivity of different crops has been studied by

many authors (Phirek et al., 1996; Pietruszewski, 1999 a, b and c and Aladjadjiyan, 2002). It has been established that the proper combination of magnetic field induction and exposure accelerates the early stages of plant development and improves the productivity. Consequently, the magnetic field effect can be used as an alternative to the chemical methods of plant treatment for improving the production efficiency Aladjadjiyan (2003). Investigations of MF on biological systems have demonstrated generalized increases in gene transcription and changes in the levels of specific mRNAs (Celik et al., 2008).

MATERIALS AND METHODS:

Two pot experiments were conducted under green house condition at Agronomy Department, National Research Centre, Dokki, Giza, Egypt during one successive winter season to evaluate magnetized water using monocotyledonous crops such as wheat (var. Sakha-93) and flax (var. Sakha-2) and dicotyledonous crops such as chick pea (var. Giza-4) and lentil (var. Sena-1).

Grains of the respective crops were obtained from Field Crop Research Department, Field Crops Institute, Agriculture Research Centre, Giza, Egypt. Grains of each crop without visible defect, insect damage and malformation were selected and planted in ten pots (30 cm in diameter and 50 cm depth) containing a mixture of clay and sandy soil [2:1]. Half of the pots were irrigated twice on a week interval with tap water, while the other 5 pots were irrigated with the tap water after magnetization through a one inch Magnetron [U.T. 3, Magnetic Technologies LLC PO Box 27559, Dubai, UAE]. The recommended NPK fertilizers for each crop were applied through the period of experiment.

After 60 days from sowing plant height, fresh and oven dry weight of 6 plants from each crop were determined. Photosynthetic pigments such as chlorophyll a, chlorophyll b and carotenoids) of leaves were estimated spectrophotometrically as the method described by Moran (1982). Total indole acetic acid (IAA) (Larsen et al., 1962), and total phenol (Malik and Singh, 1980), were estimated in the fresh leaves. Electrophoresis protein profile of leaves was analyzed according to sodium dodecylsulphate polyacrylamide gel electrophoresis [SDS-PAGE] technique (Sheri, et al., 2000). Molecular protein markers, percentage of band intensity and molecular weight of each polypeptide were related to standard markers using gel protein analyzer version 3 [MEDIA CYBERNETICS, USA].

Statistical analysis was carried out using SPSS program Version 16. A student test (Independent *t*-test) was also carried out to find the significant differences between magnetic and nonmagnetic water treatments.

RESULTS AND DISCUSSION

Growth parameters

The growth characters such as plant height, fresh and dry weight of the plants exposed to magnetic field are shown in Table 1. It is obvious that, magnetic treatment increased the growth (plant height, fresh and dry weights/plant and water content) significantly over the untreated control plants in both monocotyledonous and dicotyledonous plants. Over two seasons, the percent of increments in fresh weight/plant of monocotyledonous plants reached to 17.06 and 48.36% and reached to 10.48 and 39.25% in dry weight/plant of wheat and flax plants, respectively (Figure 1). The percent of increments in fresh and dry weight/plant of dicotyledonous plants reached to 12.51 – 18.18% and 5.76 – 15.05% in chick pea and lentil plants, respectively as compared with plants irrigated with tap water (Figure 1). Water content was found to be least affected parameters in both crops where the percent of increase ranged between 1.13 – 2.07% in all four crops. It is worthy to mention that, the percent of increases in growth parameters which reflected in fresh and dry weight/plant in this study showed that,

monocotyledonous plants (wheat and flax) surpassed dicotyledonous plants (chick pea and lentil) in their response to irrigation with magnetized.



Figure 1. Stimulation percentage over control in some growth parameters of monocot and dicot crops at 60 days after sowing.

The stimulatory effect of magnetic water may be attributed to their role in increasing absorption and assimilation of nutrients consequently increasing plant growth. These results are in good harmony with several investigators, who found that in studied paulownia tissue cultures and showed the positive effect of magnetic field on regeneration percentage (Yaycılı and Alikamanoğlu, 2005). Also, Alikamanoğlu et al. (2007) suggested that, magnetic water treatment improved seed inhibition, vigor and germination rate, and seedling treatment promoted NPK absorption and increased root no, stem thickness, dry weight/100 plants and tillers number. Moreover, Celik et al. (2008) and Nasher (2008) concluded that, magnetized water increased growth and consider an important factor for inducing chick pea plant growth. The stimulatory effect of MW on growth criteria of this study may be also attributed to the increase in photosynthetic pigment, endogenous promoters (IAA), total phenol (Table 2) and increase in protein biosynthesis (Table 4). In this connection, Shabrangi and Majd (2009) concluded that, biomass increasing needs metabolic changes particularly increasing protein biosynthesis.

Chemical constituents

Photosynthetic pigments (chlorophyll a, chlorophyll b, total chlorophyll a+b and carotenoids), total phenols and IAA contents in plant shoots exhibited remarkable increase in response to the irrigation with magnetized water when compared to untreated control plants (plants irrigated with tap water) as shown in Table 2 and Figures 2 and 3. Over of both seasons, the stimulation percent over control reached by 17.60 and 17.46% in chlorophyll a, 11.37 and 67.80% in chlorophyll b and 3.03 and 8.55 in carotenoids for monocotyledon plants (wheat and flax, respectively) and reached by 26.57 and 13.58%, 21.82 and 44.67, 0.45 and 2.70% for the above mentioned characters in dicot (chick- pea and lentil), respectively (Figure 2). Total phenol content was increased by 18.2 – 33.59% in monocot and by 20.0 – 39.02% in dicot, respectively (Figure 3). The results also showed that, total indole acetic acid content of monocot plants irrigated with magnetic water was increased by 33.35 – 233.5%, while their content in dicot plants was increased by 8.66 – 148.19% (Figure 3).

Table 1. Response of monocot and dicot crops growth at 60 days after sowing for irrigation with magnetic and normal water under green house conditions.

Treatment	Monocotyledons crops											
	Wheat						Flax					
	2008/09 season		<i>t</i> -sign.	2009/2010 season		<i>t</i> -sign.	2008/09 season		<i>t</i> -sign.	2009/2010 season		<i>t</i> -sign.
Character	Tap water	Magnetic water		Tap water	Magnetic water		Tap water	Magnetic water		Tap water	Magnetic water	
Plant height (cm)	20.75	24.12	**	26.20	29.20	**	24.00	25.00	ns	26.20	28.26	**
Fresh weight (g plant ⁻¹)	0.68	0.98	**	0.79	1.21	**	0.61	0.71	**	0.79	0.93	**
Dry weight (g plant ⁻¹)	0.17	0.23	**	0.21	0.29	**	0.15	0.16	ns	0.21	0.24	ns
Water contents (%)	75.00	76.53	Ns	74.04	75.60	ns	75.47	77.09	ns	74.10	74.19	ns
Dicotyledonous crops												
Lentil						Chick-pea						
Plant height (cm)	15.20	18.40	*	17.16	21.00	**	20.40	23.60	*	24.20	26.20	ns
Fresh weight (g plant ⁻¹)	0.56	0.66	**	0.67	0.79	*	1.39	1.58	**	1.55	1.73	**
Dry weight (g plant ⁻¹)	0.17	0.19	**	0.24	0.27	ns	0.32	0.35	ns	0.37	0.38	*
Water contents (%)	70.12	70.61	Ns	64.18	65.49	ns	76.98	77.85	ns	75.93	77.93	**

*, ** *t* is Significant at the 0.05 and 0.01 levels, respectively, ns: non significant.

Table 2. Effect of magnetic water on chemical constituents of some monocot and dicot plants at 60 days after sowing.

Treatment	Monocotyledons crops							Dicotyledons crops					
	Wheat			Flax				Chick-pea			Lentil		
Character	Tap Water	Magnetic water	<i>t</i> -sign.	Tap water	Magnetic Water	<i>t</i> -sign.	Tap water	Magnetic water	<i>t</i> -sign.	Tap water	Magnetic water	<i>t</i> -sign.	
Photosynthetic pigments (mg 100 g fresh weight ⁻¹)	Chlorophyll a	8.23	9.68	**	6.13	7.20	**	5.72	7.24	**	3.71	4.21	*
	Chlorophyll b	4.97	5.54	ns	2.36	3.96	**	3.07	3.74	**	1.25	1.80	*
	Chlorophyll a+b	13.21	15.22	**	8.49	11.16	**	8.79	10.98	**	4.96	6.02	*
	Caroteneiods	5.67	5.84	ns	4.60	4.99	ns	4.48	4.50	Ns	4.77	4.90	ns
	Total pigments	26.42	30.45	**	16.98	22.32	**	13.27	15.48	**	9.92	12.04	ns
Total phenol (mg 100 g fresh weight ⁻¹)	215.62	288.05	**	208.19	246.07	**	312.29	434.13	**	179.18	215.02	**	
Total indols (µg 100 g fresh weight ⁻¹)	2.94	9.80	**	1.19	1.59	**	1.26	1.37	**	0.83	2.05	**	

*, ** *t* is Significant at the 0.05 and 0.01 levels, respectively, ns: non significant




Figure 2. Stimulation percentage over control in photosynthetic pigments (mg 100 g fresh weight⁻¹) of monocot and dicot crops at 60 days after sowing.




Figure 3. Stimulation percentage over control in total phenols and total indoles of monocot and dicot crops at 60 days after sowing.

These results may be due to the effect of MT on alteration the key of cellular processes such as gene transcription which play an important role in altering cellular processes. It also may be due to the increase in growth promoters (Table 2). The same result was obtained by Tian et al. (1991) and Atak et al. (2000) who found an increase in chlorophyll contents specifically appeared after exposure to a magnetic field for a short time. Moreover, Atak et al. (2003) suggested that, increase all photosynthetic pigments through the increase in cytokinin synthesis which induced by MF. They also added cytokinin play an important role on chloroplast development, shoot formation, axillary bud growth, and induction of number of genes involved in chloroplast development nutrient metabolism. Atak et al. (2003) showed that, the increase in shoot regeneration, chloroplast rate, root formation and fresh weight were accompanied by the increase in auxin synthesis which induced by MF treatment of soybean plants. Moreover, Goodman et al. (1995) and Atak et al. (2003) described the role of MF in changing the characteristics of cell membrane, affecting the cell reproduction and causing some changes in cell metabolism. So the increase in total phenol under this study may be attributed to the role of MT in changing the cell membrane properties. Also, Carimi et al. (2002) and Celik et al. (2008) conclude that, MF stimulates protein synthesis via increase cytokinins and auxins and they can promote the maturation of chloroplast. Moreover, magnetic fields are known to induce biochemical changes and could be used as a stimulator for growth related reactions including affecting photosynthetic pigments (Dhawi and Al-Khayri, 2009).

Protein electrophoretic pattern

The changes in protein electrophoretic pattern of plant leaves treated with magnetic water is analyzed and recorded in Table 3. In the control leaves the separation of 12, 13, 15 and 11 protein bands were appeared in wheat, flax, chick pea and lentil, respectively. Their molecular weights ranged between 346 K Da. and 20 K Da. Magnetic water treatment of plants showed an increase in the number of protein bands to 16, 21, 22 and 16 bands in wheat, flax, chick pea and lentil, respectively. These results indicate that the leaves of plants treated with magnetic water characterized by disappearance of certain bands and the appearance of new ones as compared with that of the control plant (Table 3). The six new protein bands appeared in wheat at molecular weights 340, 194, 116, 88, 57 and 22 KDa. The nine new protein bands appeared in flax at molecular weights 301, 267, 223, 210, 113, 107, 98, 59 and 45 KDa. Also, the new protein bands appeared in chick pea at molecular weights 314, 248, 235, 226, 192, 135, 49 and 32 KDa. While in lentil, the new protein bands appeared at molecular weights 332, 307, 301, 93, 75, 55 and 38 KDa.

On the other hand, the protein bands at molecular weights 51 and 37 K Da in wheat, at 56 K KDa in chick pea and at 127 and 20 K Da. In lentil were disappeared in response to magnetic water treatment. The induction of new protein bands in response to MWT may be as a result of the effect of MFs in increases proliferation, gene expression and protein biosynthesis (Tenford, 1996). Also, Celik *et al.* (2008) found that the increase in the percentage of plant regeneration is due to the effect of MF of cell division and protein synthesis in paulownia node cultures, and concluded that, investigations of MF on biological systems have demonstrated generalized increases in gene transcription and changes in the levels of specific mRNAs. Moreover, Shabrangi and Majd (2009) concluded that, biomass increasing needs metabolic changes particularly increasing protein biosynthesis. They also add magnetic field is known as an environmental factor which affects on gene expression. Therefore, by augmentation of biological reactions like protein synthesis, biomass would increase too.

Yield and yield component

With respect to the effect of MT on the yield and yield component of monocot and dicot plants data in Tables 4a and 4b cleared that MT increased all yield characters in all crops over the untreated controls. Over two seasons, the percentage of increase reached to 3.89 and 11.88% in plant height, 3.64 and 6.72% in 100-seed weight and 9.10 and 31.33% in seed yield/plant for monocotyledonous crops (flax and wheat, respectively) and to 17.98 and 26.48%, 6.24 and 2.54% and to 24.98 and 38.64% in above mentioned parameters for dicotyledonous crops (lentil and chick pea, respectively) over the untreated controls (Figure 4).

Generally, the stimulatory effect of magnetic treatment may be attributed to their role in increasing growth (Table 1), photosynthetic pigment and growth promoters (Table 2) consequently increasing yield characters. These results are in agreement with those obtained by Tian *et al.* (1991) who indicated that, MW increased yield of rice by 13.2%. This was accompanied the stimulation effect of MW on leaf chlorophyll content. Kordas (2002) found that, the exposure of green tops and root systems of wheat plant to MF increased quantity of coarse grain by 10.6% and 6.3% respectively. In this connection, Dodlesny *et al.* (2004, 2005) suggested that, the gain in seed yield resulting from the pre-sowing treatment of seeds with MF for broad bean and pea was due to the higher number of pods per plant and the fewer plant losses in the unit area in the growing season. Moreover, Souza *et al.* (2006) showed that, MT on tomato increased significantly the mean fruit weight, the fruit yield/plant, the fruit yield per area and the equatorial diameter of fruits in comparison with the controls. Moreover, MF was shown to induce fruit yield per plant and average fruit weight (Celik *et al.*, 2008). Exposure of plants to MW is highly effective in enhancing growth characteristics. This observation suggests that there may be resonance-like phenomena which increase the internal energy of the seed that occurs. Therefore, it may be possible to get higher yield (Vashisth *et al.*, 2008 and Shabrangi and Majd,

2009) on chickpea and lentil respectively.

Table 3. The relative area percentage of protein bands in leaves at 60 days after sowing of some monocot and dicot plants irrigated with magnetized and normal water

M wt. K.Da.	Monocot				Dicot			
	Wheat		Flax		Chickpea		Lentil	
342	4.54	2.24					1.76	1.83
339		5.28	2.54	3.58				
327	5.07	4.16			3.54	2.51		2.44
323					3.61	1.47	2.72	2.58
322					16.97	1.46		
316			4.49	3.03		2.32		
307				3.42				2.00
301								1.27
286	8.73	8.21	2.43	3.03				
267				3.22				
253	8.21	7.34	2.37	3.26				
245						7.80	10.23	12.75
233			4.61	3.15		1.31		
224				3.65		2.51		
204		10.32		2.40		2.68		
189					2.32	2.32		
146	13.79	6.80	7.40	7.60	4.23	5.38	8.60	4.97
135						1.99	4.55	2.15
125					2.06	2.06	6.17	
114		6.11		3.86				
107				2.56				
95				5.49				10.35
87		4.55	15.70		4.27	5.24		
73			6.21	2.75				7.85
66	9.67	9.81			6.89	7.70	17.32	2.85
56		3.92		6.77	6.15			6.05
52	8.08		10.61	6.03	2.76	5.69	8.11	4.73
47			11.22	10.44		5.08	13.30	14.96
45	15.47	12.84		5.96	10.45	7.76		
41			11.36	4.75	8.80	7.17		
37	5.16		7.26	2.45	8.79	6.15		6.59
35					5.34	2.25		
33	11.67	9.23	13.80	12.60		10.68	18.31	16.63
30					5.60	8.47		
22		4.24						
20	9.61	4.95					8.93	
Band number	11	15	13	21	15	22	11	16
Number of new band		6		9		8		7

CONCLUSION

In summary, growth parameters, biochemical components and yield components of tested plants is concomitantly increased when plants are treated with magnetic water. Biochemical components such as photosynthetic pigments, endogenous total indole; total phenol and protein contents were also increased significantly in plants treated with magnetic water. The variation in the response of plants should need continuous efforts from researchers to explore the mode of magnetic treatment action in monocot and dicot crops.

Table 4a. Response of wheat and flax yield and its components at harvest for irrigation with magnetic and normal water.

Treatment	Wheat					<i>t</i> -sign.
	2008/09 season		<i>t</i> -sign.	2009/2010 season		
Character	Tap water	Magnetic water			Tap water	Magnetic water
Plant height (cm)	39.80	47.00	*	56.40	59.60	*
Spike length (cm)	5.00	6.60	**	8.50	9.20	**
Spike weight (g)	0.48	0.53	**	0.64	0.75	**
Spikeletes no. spike ⁻¹	9.00	12.00	**	14.40	16.00	**
100-grain weight (g)	4.04	4.31	ns	4.14	4.42	ns
Grain yield (g tiller ⁻¹)	0.30	0.40	**	0.75	0.97	**
Straw yield (g tiller ⁻¹)	0.59	0.80	**	0.93	1.06	**
Biological yield (g tiller ⁻¹)	0.89	1.20	**	1.68	2.03	**
HI (%)	33.63	33.33	ns	44.64	47.78	ns
	Flax					
Plant height (cm)	56.80	58.20	*	58.30	61.40	**
Tecenical length (cm)	43.40	48.80	*	48.50	51.60	**
Based branches (number plant ⁻¹)	2.40	2.80	ns	2.60	2.84	ns
Fruit Branches (number plant ⁻¹)	5.60	6.00	ns	6.20	6.44	ns
Cabsules (number plant ⁻¹)	9.20	10.80	ns	10.40	11.60	ns
Cabsules weight (g plant ⁻¹)	0.44	0.53	*	0.53	0.57	ns
Seed (number cabsula ⁻¹)	8.00	8.40	ns	8.26	9.28	**
Seeds (number plant ⁻¹)	73.60	90.72	**	85.68	107.46	**
100-seed weight (g)	0.68	0.70	ns	0.69	0.72	ns
Seed yield (g plant ⁻¹)	0.32	0.35	*	0.34	0.37	ns

*, ** *t* is Significant at the 0.05 and 0.01 levels, respectively, ns: non significant.




Figure 4. Stimulation percentage over control in some yield parameters of monocot and dicot crops at 60 days after sowing.

ACKNOWLEDGEMENTS

We are most appreciation of honorable Prof. P.Ponmurugan, Head, Dept. of Biotechnology K.S.R. College of Technology Tiruchengode - 637 215 Namakkal Dist, Tamil Nadu, India and Prof. Abd El-Lateef E.M., National Research Centre for revision this manuscript

Table 4 b. Response of lentil and chick-pea yield and its components at harvest for irrigation with magnetic and normal water.

Treatment	Lentil						Chick-pea					
	2008/09 season		<i>t</i> -sign.	2009/2010 season		<i>t</i> -sign.	2008/09 season		<i>t</i> -sign.	2009/2010 season		<i>t</i> -sign.
Character	Tap water	Magnetic water		Tap water	Magnetic water		Tap water	Magnetic water		Tap water	Magnetic water	
Plant height (cm)	16.40	20.60	**	23.20	25.60	*	28.40	35.20	**	32.40	41.80	**
Branches (number plant ⁻¹)	2.71	3.60	*	3.32	3.92	*	2.47	3.23	**	3.20	4.40	**
Pods (number plant ⁻¹)	4.78	6.40	**	6.76	8.40	**	6.60	8.81	ns	7.60	11.50	**
Pods weight (g plant ⁻¹)	0.63	0.72	*	0.74	0.88	**	1.86	2.59	**	1.96	2.76	**
Seeds (number plant ⁻¹)	8.75	10.50	**	10.66	12.34	**	6.89	9.50	**	7.13	10.20	**
100-seed weight (g)	5.20	5.62	**	5.45	5.69	**	18.16	19.03	**	19.13	19.17	ns
Seed yield (g plant ⁻¹)	0.52	0.66	**	0.63	0.78	**	1.36	1.77	**	1.43	2.10	**
Straw yield (g plant ⁻¹)	0.54	0.71	**	0.75	0.91	*	1.43	1.91	**	1.98	2.94	**
Biological yield (g plant ⁻¹)	1.06	1.37	**	1.38	1.69	**	2.79	3.68	**	3.41	5.04	**

*, ** *t* is significant at the 0.05 and 0.01 levels, respectively, ns: non significant.

REFERENCES

- Aladadjjiyan A. 2002. Study of the Influence of Magnetic Field on Some Biological Characteristics of *Zea mais*. J. of Central Europ. Agric. 3(2). 89-94.
- Aladadjjiyan A. 2003. Use of physical factors as an alternative to chemical amelioration. J. Environ. Protec. and Ecol. 4 (3): 662-667.
- Aladadjjiyan A, Ylieva T 2003. Influence of stationary magnetic field on the early stages of the development of tobacco seeds (*Nicotiana tabacum* L.). Journal of Central European Agriculture 132 ,4 (2): 131-138.
- Alikamanoğlu S, Yaycılı O, Atak Ç, Rzakoulieva A. 2007. Effect of magnetic field and gamma radiation on paulownia tomentosa tissue culture. J. Biotechnology & Biotechnological Equipment 21 (1): 21:49-53
- Atak C, Danilov V, Yurttas B, Yalçın S, Mutlu D, Rzakoulieva A. 1997. Effects of magnetic field on soybean (*Glycine max* L.Merrill) seeds. Com JINR. Dubna 1-13.
- Atak Ç, Danilov V, Yurttas B, Yalçın S, Mutlu D, Rzakoulieva A. 2000. Effect of magnetic field on Paulownia seeds. Com JINR. Dubna. 1-14.
- Atak C, Emiroglu O, Aklimanoglu S, Rzakoulieva A. 2003. Stimulation of regeneration by magnetic field in soybean (*Glycine max* L. Merrill) tissue cultures. J Cell Mol. Biol. 2:113–119.
- Atak C, Celik O, Olgun A, Alikamanolu S, Rzakoulieva A. 2007. Effect of magnetic field on peroxidase activities of soybean tissue culture. Biotechnology 21: 166-71.
- Carimi F, Zottini M, Formentin E, Terzi M, Schiaw FL, 2002. Cytokinins new apoptotic inducers in plants, Planta 216 (3): 413-421.
- Çelik Ö, Atak C, Rzakoulieva A. 2008. Stimulation of rapid regeneration by a magnetic field in paulownia node cultures. Journal of Central European Agriculture 9 (2): 297 – 303.
- De Souza A, Garcí D, Sueiro L, Gilart F, Porrás E, Licea L. 2006. Pre-sowing magnetic treatments of tomato seeds increase the growth and yield of plants. Bioelectromagnetics 27, 247-257.
- Dhawi F, Al-Khayri JM. 2009. Magnetic fields induce changes in photosynthetic pigments content in date palm (*Phoenix dactylifera* L.). Seedlings the open agriculture journal 3, 1-5.
- Goodman EM, Greenbaum B, Morron TM. 1995. Effects of electromagnetic fields on molecules and cells. International Review of Cytology 158: 279-325.
- Kordas L. 2002. the effect of magnetic field on growth, development and the yield of spring wheat. Polish journal of environmental studies 11 (5): 527-530.
- Larsen P, Harbo, A, Klungron S, Ashein TA 1962. On the biosynthesis of some indole compounds in *Acetobacter xylinum*. Physiol Plant. 15: 552-565.
- Malik CP, MB Singh 1980. Plant Enzymology and Histoe-nzymology. Kalyani Publishers. New Delhi.
- Moran R. 1982. Formulae for determination of chlorophyllous pigments extracted with N, N-dimethylformamide. Plant Physiol. 69:1371-1381.
- Namba K Sasao A, Shibusawa S. 1995. Effect of magnetic field on germination and plant growth. Acta Horticulture 399: 143-147.
- Nasher SH. (2008). The Effect of magnetic water on growth of chick-pea seeds Eng. & Tech. 26,(9):4 pages.
- Oldacay S, Erdem G. 2002. Evaluation of chlorophyll contents and peroxides activities in (*Helianthus annuus* L.) genotypes exposed to radiation and magnetic field. Pakistan Journal of Applied Science 2(10): 934-937.
- Phirke PS, Kudbe AB, Umbarkar SP. 1996. The influence of magnetic field on plant growth. Seed Sci. Technol. 24, 375-392.
- Pietruszewski S. 1999 b. Influence of pre-sowing magnetic biostimulation on germination and yield of wheat. Int. Agrophysics 13, 241-244.
- Pietruszewski S. 1999 c. Magnetic treatment of spring wheat seeds (in Polish). Rozprawy NaukoweARin Lublin 220, 1-55.
- Pietruszewski ST. 1999 a. Effect of alternating magnetic field on germination, growth and yield of plant seeds. Inzynieria Rolnicza 5(11) 209-215.
- Podlesny J, Pietruszewski S, Podlesna A. 2005. Influence of magnetic stimulation of seeds on the formation of morphological features and yielding of the pea. Int Agrophys 19: 1–8.
- Podlesny J, Pietruszewski S, Podleona A. 2004. Efficiency of the magnetic treatment of broad bean seeds cultivated under experimental plot conditions. Int. Agrophysics 18, 65 - 71.
- Reina FG, Pascual LA, Fundora IA. 2001. Influence of a Stationary Magnetic Field on Water Relations in Lettuce Seeds. Part II: Experimental Results Bioelectromagnetics 22:596-602.

- Shabrangi A, Majd A. 2009. Effect of Magnetic Fields on Growth and Antioxidant Systems in Agricultural Plants PIERS Proceedings, Beijing, China, March 23-27.
- Sheri LH, Ncolas ES, Michae TK Joanna BG. 2000. Comparison of protein expressed by *Pseudomonas aeruginosa* strains representing initial and chronic isolates from a cystic fibrosis patient: an analysis by 2 – D gel electrophoresis and capillary coloumn liquid chromatograph tandem mass spectrometry. *Microbiology* 146: 2495 – 2508.
- Tenforde TS 1996. Interaction of ELF magnetic field with living systems. In: Handbook of biological effects of electromagnetic fields. Polk C, Postow E. (Ed). Second Edition, CRC Press USA 85-230.
- Tian WX, Kuang YL, Mei ZP (1991). Effect of magnetic water on seed germination, seedling growth and grain yield of rice. *Field Crop Abstracts*. 044-07228.
- Vashisth A. and Nagarajan, S. 2008. Exposure of seeds to static magnetic field enhances germination and early growth characteristics in chickpea (*Cicer arietinum* L.). *Bioelectromagnetics* 29:571-578.
- Wo'jcik S. 1995. Effect of the pre-sowing magnetic biostimulation of the Buckwheat seeds on the yield and chemical composition of Buckwheat grain. *Current Adv. uckwheat Res* 93:667– 674.
- Yaycılı O, Alikamanoğlu S. 2005. The effect of magnetic field on Paulownia tissue cultures. *Plant Cell Tissue and Organ Culture* 83 (1): 1109-114.